


Sexually Transmitted Disease in Milwaukee County

Wisconsin STD Surveillance Data
Cases diagnosed in 2006



Reported STD case rates*, by county, reportable STDs**, Wisconsin, 2006



Counties with rates* >400


<u>County</u>	<u>Cases</u>	<u>Percent</u>	<u>Rate*</u>
Milwaukee	16,004	52.7%	1,712
Menominee	65	0.2%	1,412
Racine	1,481	4.9%	768
Forest	56	0.2%	553
Kenosha	822	2.7%	522
Rock	811	2.7%	519
Dane	2,339	7.7%	516
Brown	1,113	3.7%	466
La Crosse	471	1.5%	430
Dodge	363	1.2%	410
Subtotal	23,525	77.4%	1,003
Other counties	6,870	22.6%	215
State total	30,395	100.0%	549

* Cases per 100,000 population.

** Chlamydia, gonorrhea, syphilis and genital HSV infection.



Reported STD cases, Milwaukee County, 2006




<u>Disease</u>	<u>Cases</u>	<u>Rate*</u>
Chlamydia	9,975	1,066
Gonorrhea	4,896	523
Syphilis	90	9
HSV	1,043	111
All STDs	16,004	1,712

* Cases per 100,000 population.




Primary and secondary syphilis rates* for the 50 largest metropolitan areas, 2005, CDC data



* Cases per 100,000 population.




Chlamydia rates* for the 50 largest metropolitan areas, 2005, CDC data




* Cases per 100,000 population.

Gonorrhea rates* for the 50 largest metropolitan areas, 2005, CDC data



* Cases per 100,000 population.

Combined rates* for chlamydia, gonorrhea and primary and secondary syphilis for the 50 largest metropolitan areas, 2005, CDC data





* Cases per 100,000 population.



Reported STD rates* by zip code, reportable STDs**, Milwaukee County, 2006

Rate*



Zip codes with rates* >3,000

<u>Zip code</u>	<u>Cases**</u>	<u>Percent</u>	<u>Rate*</u>
53203	25	<1%	7,463
53206	1,815	11%	5,522
53205	550	3%	5,057
53210	1,488	9%	4,877
53212	1,153	7%	3,743
53216	1,208	8%	3,566
53208	1,251	8%	3,559
53218	1,378	9%	3,401
53209	1,533	10%	3,127
Subtotal	10,401	65%	3,941
Other zip code	5,603	35%	835
Milwaukee Co.	16,004	100%	1,712

* Cases per 100,000 population.

** Chlamydia, gonorrhea, syphilis and genital HSV infection.

