


Ending
Conversion Therapy:
Supporting and Affirming
LGBTQ Youth



October 2015



PAGE INTENTIONALLY LEFT BLANK



Ending Conversion Therapy:
Supporting and Affirming
LGBTQ Youth

October 2015





PAGE INTENTIONALLY LEFT BLANK



Acknowledgements

This report was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA) by Abt Associates under contract number HHSS283200700008I/HHSS28342001T with SAMHSA, U.S. Department of Health and Human Services (HHS). David Lamont Wilson served as the Government Project Officer. Elliot Kennedy served as the Task Lead.

Disclaimer

The views, opinions, and content of this publication are those of the author and do not necessarily reflect the views, opinions, or policies of SAMHSA or HHS. Listings of any non-Federal resources are not all-inclusive and inclusion of a listing does not constitute endorsement by SAMHSA or HHS.

Public Domain Notice

All material appearing in this report is in the public domain and may be reproduced or copied without permission from SAMHSA. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Electronic Access and Printed Copies


This publication may be downloaded or ordered at <http://store.samhsa.gov>. Or call SAMHSA at 1-877-SAMHSA-7 (1-877-726-4727) (English and Español).

Recommended Citation

Substance Abuse and Mental Health Services Administration, Ending Conversion Therapy: Supporting and Affirming LGBTQ Youth. HHS Publication No. (SMA) 15-4928. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2015.


Originating Office

Division of Systems Development, Center for Substance Abuse Prevention, Substance Abuse and Mental Health Services.



Contents

Executive Summary	1
Introduction	7
Professional Consensus Process	9
Statements of Professional Consensus.....	11
Research Overview.....	15
Approaches to Ending the Use of Conversion Therapy	37
Guidance for Families, Providers, and Educators	41
Summary and Conclusion.....	51
References.....	52
Appendix A: Glossary of Terms	64
Appendix B: Acknowledgments	64
Endnotes	66



Executive Summary

Lesbian, gay, bisexual, and transgender youth, and those who are *questioning* their sexual orientation or gender identity (*LGBTQ* youth) experience significant health and behavioral health disparities. Negative social attitudes and discrimination related to an individual's *LGBTQ* identity can contribute to these disparities, and may result in institutional, interpersonal, and individual stressors that affect mental health and well-being. (Bockting, Miner, Swinburne Romine, Hamilton, & Coleman, 2013; Meyer, 2003). This stress, as well as limited opportunities for support, are encountered by *sexual and gender minority* youth in their families, communities, and school settings. Additionally, some transgender youth experience gender dysphoria – psychological distress due to the incongruence between one's body and gender identity (Coleman et al., 2012).

SAMHSA is committed to eliminating health disparities facing vulnerable communities, including sexual and gender minority communities. One key factor to preventing these adverse outcomes is positive family (including guardians and caregivers) and community engagement and appropriate interventions by medical and behavioral health care providers. Supporting optimal development of children and adolescents with regard to sexual orientation, gender identity, and gender expression is vital to ensuring their health and well-being.

The purpose of this report, *Ending Conversion Therapy: Supporting and Affirming LGBTQ Youth*, is to provide mental health professionals and families with accurate information about effective and ineffective therapeutic practices related to children's and adolescent's sexual orientation and gender identity. Specifically, this report addresses the issue of conversion therapy for minors. The conclusions in this report are based on professional consensus statements arrived at by experts in the field. Specifically, conversion therapy—efforts to change an individual's sexual orientation, gender identity, or gender expression²—is a practice that is not supported by credible evidence and

has been disavowed by behavioral health experts and associations. Conversion therapy perpetuates outdated views of gender roles and identities as well as the negative stereotype that being a sexual or gender minority or identifying as *LGBTQ* is an abnormal aspect of human development. Most importantly, it may put young people at risk of serious harm.

Key Findings

This report and its recommendations are based on consensus statements developed by experts in the field after a careful review of existing research, professional health association reports and summaries, and expert clinical guidance. The consensus statements highlight areas of the ethical and scientific foundations most relevant to the practice of conversion therapy with minors. A full list of the consensus statements is found in the body of this report; key statements that form the underpinnings of the guidance in this report are provided here.

- Same-gender³sexual orientation (including identity, behavior, and attraction) and variations in gender identity and gender expression are a part of the normal spectrum of human diversity and do not constitute a mental disorder.
- There is limited research on conversion therapy efforts among children and adolescents; however, none of the existing research supports the premise that mental or behavioral health interventions can alter gender identity or sexual orientation.
- Interventions aimed at a fixed outcome, such as gender conformity or heterosexual orientation, including those aimed at changing gender identity, gender expression, and sexual orientation are coercive, can be harmful, and should not be part of behavioral health treatment. (American Psychiatric Association, 2013b; American Psychological Association, 2010; National Association of Social Workers, 2008).

Understanding Sexual Orientation and Gender Identity in Children and Youth

Behavioral health providers, parents, schools, and communities can best provide support to children, adolescents, and their families when they have access to the most current information about sexual orientation, gender identity, and gender expression in youth. The following overview presents the best current evidence regarding understandings of child and adolescent sexual orientation, gender identity, and gender expression.

Sexuality occurs across a continuum; same-gender attraction and relationships are normal variations of human sexuality (Diamond, 2015; Vrangalova & Savin-Williams, 2012). Similarly, a gender identity that is incongruent with assigned sex at birth, as well as a gender expression that diverges from stereotypical cultural norms for a particular gender, are normal variations of human gender (American Psychological Association, 2015a; Knudson, De Cuypere, & Bockting, 2010). Being a sexual or gender minority, or identifying as LGBTQ, is not pathological (American Psychological Association, 2015a; APA Task Force on Gender Identity and Gender Variance, 2009; Coleman et al., 2012).

There is not a single developmental trajectory for either sexual minority or gender minority youth. Compared to the 20th century, in the 21st century, youth started realizing and disclosing a minority sexual orientation and/or identifying as lesbian, gay, or bisexual at younger ages than in previous generations (Diamond & Savin-Williams, 2000; Floyd & Bakeman, 2006; Grov, Bimbi, Nanín, & Parsons, 2006; R. C. Savin-Williams, 2001). Though aspects of sexuality are displayed beginning in infancy, little is known about sexual orientation among pre-pubertal children (Adelson & American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI), 2012). Children are rarely if ever distressed about their current or future sexual orientation; more commonly, parents and guardians are distressed about a child's perceived current or future sexual orientation and seek the

assistance of behavioral health providers (American Psychological Association, 2009). Sexual minority adolescents face the same developmental tasks that accompany adolescence for all youth, including sexual orientation identity development. Unlike those with a heterosexual orientation, however, adolescents with a minority sexual orientation must navigate awareness and acceptance of a socially marginalized sexual identity; potentially without family, community, or societal support. In comparison with their heterosexual counterparts, sexual minority adolescents are at increased risk for psychological distress and substance use behaviors, including depressive symptoms, increased rates of substance use and abuse, suicidal ideation and attempts, as well as increased likelihood of experiencing victimization, violence, and homelessness (Corliss et al., 2010; Friedman et al., 2011; Goldbach, Tanner-Smith, Bagwell, & Dunlap, 2014; Hatzenbuehler, 2011; Institute of Medicine, 2011; Kann et al., 2011; Marshal et al., 2011; Russell, 2003). Supportive families, peers, and school and community environments are associated with improved psychosocial outcomes for sexual minority youth (Bouris et al., 2010; Kosciw, Greytak, Palmer, & Boesen, 2014; Lease, Horne, & Noffsinger-Frazier, 2005).

Gender development begins in infancy and continues progressively throughout childhood. Gender diversity or signs of gender dysphoria may emerge as early as a child's preschool years, or as late as adolescence (Cohen-Kettenis, 2005). For many gender minority children, gender dysphoria will not persist, and they will develop a *cisgender* identity in adolescence or adulthood; a majority of these children will identify as lesbian, gay, or bisexual in adulthood (Bailey & Zucker, 1995; Drescher, 2014; Leibowitz & Spack, 2011; Wallien & Cohen-Kettenis, 2008). Whether or not these individuals continue to have a diverse gender expression is unknown. For other gender minority children, gender dysphoria will persist and usually worsen with the physical changes of adolescence; these youth generally identify as transgender (or another gender identity that differs from their assigned sex at birth) in adolescence and adulthood

(Byne et al., 2012; Coleman, et al., 2012). For still another group, gender dysphoria emerges in post-puberty without any childhood history of gender dysphoria gender diversity (Edwards-Leeper & Spack, 2012). Gender dysphoria that worsens with the onset of puberty is unlikely to remit later in adolescence or adulthood, especially among youth with a childhood onset, and long-term identification as transgender is likely (American Psychological Association, 2015a; American Psychological Association, 2008; Byne, et al., 2012).

While most adolescents with gender dysphoria score within normal ranges on psychological tests (Cohen-Kettenis & van Goozen, 1997; de Vries, Doreleijers, Steensma, & Cohen-Kettenis, 2011; Smith, van Goozen, & Cohen-Kettenis, 2001), some gender minority children and adolescents have elevated risk of depression, anxiety, and behavioral issues. These psychosocial issues are likely related to if not caused by negative social attitudes or rejection (Vance, Ehrensaft, & Rosenthal, 2014). As with sexual minority adolescents, other issues of clinical relevance for gender minority adolescents include increased risk of experiencing victimization and violence, suicidal ideation and attempts, and homelessness (Coleman, et al., 2012; Garofalo, Deleon, Osmer, Doll, & Harper, 2006; Institute of Medicine, 2011; Mustanski, Garofalo, & Emerson, 2010; Simons, Leibowitz, & Hidalgo, 2014). Improved psychosocial outcomes are seen among youth when social supports are put in place to recognize and affirm gender minority youth's gender identities (Vance, et al., 2014).

Therapeutic Efforts with Sexual and Gender Minority Youth⁴

Given the professional consensus that conversion therapy efforts are inappropriate, the following behavioral health approaches are consistent with the expert consensus statements and current research, and are recommended by professional associations (American Psychological Association, 2015a; APA Task Force on Appropriate Therapeutic Responses to Sexual Orientation, 2009; Byne, et al., 2012). When providing services to children, adolescents, and families, appropriate therapeutic approaches

include: providing accurate information on the development of sexual orientation and gender identity and expression; increasing family and school support; and reducing family, community, and social rejection of sexual and gender minority children and adolescents. Approaches should be client-centered and developmentally-appropriate with the goal of treatment being the best possible level of psychological functioning, rather than any specific gender identity, gender expression, or sexual orientation. Appropriate therapeutic approaches with sexual and gender minority youth should include a comprehensive evaluation and focus on identity development and exploration that allows the child or adolescent the freedom of self-discovery within a context of acceptance and support. It is important to identify the sources of any distress experienced by sexual and gender minority youth and their families, and work to reduce this distress. Working with parents and guardians is important as parental behaviors and attitudes have a significant effect on the mental health and well-being of sexual and gender minority children and adolescents. School and community interventions may also be necessary and appropriate.

In addition to the appropriate therapeutic approaches described above – comprehensive evaluation, support in identity exploration and development without an *a priori* goal of any particular gender identity or expression, and facilitation of family and community support – social transition and medical intervention are therapeutic approaches that are appropriate for some gender minority youth. Careful evaluation and developmentally-appropriate informed consent of youth and their families, including a weighing of potential risks and benefits are vital when considering medical intervention with gender minority youth.

Eliminating the practice of conversion therapy with sexual and gender minority minors is an important step, but it will not alleviate the myriad of stressors they experience as a result of interpersonal, institutional, and societal bias and discrimination against sexual and gender minorities.

LGBTQ youth still need additional support to promote positive development in the face of such stressors. Supportive family, community, school, and health care environments have been shown to have great positive impacts on both the short- and long-term health and well-being of LGBTQ youth. Families and others working with LGBTQ children and adolescents can benefit from guidance and resources to increase support for sexual and gender minority minors and to help facilitate the best possible outcomes for these youth.

Ending the Use of Conversion Therapy for Minors

Given that conversion therapy is not an appropriate therapeutic intervention; efforts should be taken to end the practice of conversion therapy. Efforts to end the practice have included policy efforts to reduce the negative attitudes and discrimination directed at LGBTQ individuals and families; affirmative public information about LGBTQ individuals, particularly directed at families and youth; resolutions and guidelines by professional associations to inform providers that conversion efforts are inappropriate and to provide guidance on appropriate interventions; and, state and federal legislation and legal action to end the practice of conversion therapy. Future efforts may include improved provider training, federal regulatory action, advancement of legislation at the state and federal level, and additional activities by the Administration, which issued a public statement supporting efforts to ban the use of conversion therapy for minors in the spring of 2015.

PAGE INTENTIONALLY LEFT BLANK



Introduction

This report, *Ending Conversion Therapy: Supporting and Affirming LGBTQ Youth*, provides an overview of the current state of scientific understanding of the development of sexual orientation and gender identity in children and adolescents as well as the professional consensus on clinical best practices with these populations. Specifically, this report addresses the issue of conversion therapy for minors. Conversion therapy—efforts to change an individual’s sexual orientation, gender identity, or gender expression⁵—is a practice that is not supported by credible evidence, and has been disavowed by behavioral health experts and associations. Importantly, this report also provides a nuanced overview of appropriate supportive interventions to assist families in exploring the sometimes difficult issues associated with sexual orientation, gender identity, and gender expression.

This work is the result of a collaboration between the Substance Abuse and Mental Health Services Administration (SAMHSA) and the American Psychological Association (APA), which convened a panel of behavioral health professionals (e.g., psychologists, researchers and clinicians from psychology, social work, and psychiatry) with expertise in the fields of gender development, gender identity, and sexual orientation in children and adolescents in July 2015. That convening, which is discussed in greater depth below, aimed to establish consensus with respect to conversion therapy for minors, based on the best available research and scholarly material available, as well as the clinical experience of experts in the field. The resultant statements of professional consensus are printed in their entirety in the following section.

In addition, this report highlights [areas of opportunity for future research](#), and provides an overview of [mechanisms to eliminate the use of harmful therapies](#). In an effort to provide useful tools for families, practitioners, and educators, the report also provides resources on several topics, including: [Family and Community Acceptance](#),


“Being gay is not a disorder. Being transgender is not a malady that requires a cure.”

—Vice Admiral Vivek H. Murthy,
19th U.S. Surgeon General

[School-Based Issues](#), [Pediatric Considerations](#), and [Affirmative Exploratory Therapies](#). In addressing these four topics, SAMHSA aims to enable families, providers, educators, and community members to take actions that will reduce the health risks and disparities facing this vulnerable population.

SAMHSA is committed to eliminating health disparities facing vulnerable communities, including sexual and gender minority communities. In addressing the issues included in this report that have a significant impact on the lives and well-being of sexual and gender minority youth, SAMHSA aims to enable families, providers, and educators to take actions that will reduce the health risks and disparities facing this vulnerable population.

SAMHSA’s mission is to improve the behavioral health of the nation. As such, SAMHSA endeavors to improve public health and eliminate health disparities facing all vulnerable communities, including sexual and gender minority populations.⁶As will be addressed in detail below, conversion therapy perpetuates outdated gender roles and negative stereotypes that being a sexual or gender minority or identifying as LGBTQ is an abnormal aspect of human development. Most importantly, it may put young people at risk of serious harm. This report is one of many steps SAMHSA is taking to improve the health and well-being of sexual and gender minority children and youth.



Professional Consensus Process

In early April 2015, representatives from SAMHSA and APA agreed to collaborate to address the concerns of professional associations, policy makers, and the public regarding efforts to change gender identity and sexual orientation in children and adolescents (also referred to as conversion therapy). Through the support of the Federal Agencies Project, APA hosted an expert consensus convening on this topic in July 2015, which significantly informed this report. The research overview and clinical expertise highlighted throughout serve as the foundation from which the consensus statements were developed. Both the process of achieving consensus and the results of the meeting are published below.


APA initially developed a list of the areas of expertise to be used in identifying potential experts to participate in the consensus panel based on existing professional guidelines and resolutions related to sexual orientation, gender identity, and gender expression, as well as published research. APA solicited nominations from specialists in the field with expertise in gender, sexuality and sexual orientation, child and adolescent development and mental health, and the psychology of religion. Additionally, APA solicited nominations from professional associations representing the major mental health and health professions. Using the input received from these sources, APA extended invitations to a short list of highly recommended group of experts. This initial expert pool nominated additional experts based on their assessment of the expertise needed to achieve the goals of the meeting. The final panel of 13 experts consisted of ten psychologists, two social workers, and one psychiatrist. These individuals included researchers and practitioners in child and adolescent mental health with a strong background in gender development, gender identity, and sexual orientation in children and adolescents. The panel also included experts with a background in family therapy, ethics, and the psychology of religion. Among others, the panel included: Sheri Berenbaum, PhD; Celia B.

Fisher, PhD; Laura Edwards-Leeper, PhD; Marco A. Hidalgo, PhD; David Huebner, PhD; Colton L. Keo-Meier, PhD; Scott Leibowitz, MD; Robin Lin Miller, PhD; Caitlin Ryan, PhD, ACSW; Josh Wolff, PhD; and Mark Yarhouse, PsyD. APA activities were coordinated by Clinton W. Anderson, PhD and Judith Glassgold, PsyD.

Based on published literature on consensus methods, APA developed an iterative process that culminated in a two-day meeting in Washington, DC on July 7 and 8, 2015. During the meeting, panelist-led discussions considered the relevant research, professional guidelines and clinical knowledge-base for each of the topics. The panel developed consensus statements on sexual orientation change efforts as well as gender identity change efforts in children and adolescents for each of the relevant developmental stages: pre-pubertal children, peri-pubertal adolescents, and pubertal and post-pubertal adolescents.

Panelists agreed that unanimous consensus was a strong priority, but that if unanimity could not be reached, 80 percent support would constitute consensus. The panelists also agreed that minority opinions should be reflected in the record if any dissenting expert wished to issue such an opinion. Unanimous consensus was reached in nearly all instances. No dissenting opinions were formally registered. The statements of professional consensus are printed in *Section 3* of this report.

Observers from interested federal agencies, health and human services professional organizations, foundations, and LGBTQ human rights organizations also attended the meeting. These observers were offered an opportunity to submit written questions, which the panel addressed throughout the course of the meeting.



“PFR” created something that was groundbreaking. It was the best piece of work ever created by the federal government.”

Sue Thau

Statements of Professional Consensus

The following are the statements of professional consensus regarding sexual orientation and gender identity and expression that were developed during the July 2015 APA consensus convening. After initially developing separate statements regarding issues relating to the development of sexual orientation and gender identity and gender expression, the panel developed a set of three key summary statements. The panel also developed a statement regarding the guiding human rights and scientific principles that provide a foundation for behavioral health professionals' work in this area.

Guiding Principles

Behavioral health professionals respect human dignity and rights. The foundational ethical principle of “self-determination” requires that children and adolescents be supported in their right to explore, define, and articulate their own identity. The principles of “justice” and “beneficence and nonmaleficence” require that all children and adolescents have access to behavioral health treatments that will promote their health and welfare. Children and adolescents have the right to participate in decisions that affect their treatment and future. Behavioral health professionals respect human diversity and strive to incorporate multicultural awareness into their work.

These guiding principles are based upon the codes of ethics for the professional fields of Psychology, Psychiatry, and Social Work (American Psychiatric Association, 2013b; American Psychological Association, 2010; National Association of Social Workers, 2008).

Professional Consensus on Conversion Therapy with Minors

1. Same-gender⁷sexual orientation (including identity, behavior, and/or attraction) and variations in gender identity and gender expression are a part of the normal spectrum of human diversity and do not constitute a mental disorder.
2. There is limited research on conversion therapy efforts among children and adolescents; however, none of the existing research supports the premise that mental or behavioral health interventions can alter gender identity or sexual orientation.
3. Interventions aimed at a fixed outcome, such as gender conformity or heterosexual orientation, including those aimed at changing gender identity, gender expression, and sexual orientation are coercive, can be harmful, and should not be part of behavioral health treatments. Directing the child to be conforming to any gender expression or sexual orientation, or directing the parents to place pressure for specific gender expressions, gender identities, and sexual orientations are inappropriate and reinforce harmful gender and sexual orientation stereotypes.

Professional Consensus on Sexual Orientation in Youth

1. Same-gender sexual identity, behavior, and attraction are not mental disorders. Same-gender sexual attractions are part of the normal spectrum of sexual orientation. Sexual orientation change in children and adolescents should not be a goal of mental health and behavioral interventions.
2. Sexual minority children and adolescents are especially vulnerable populations with unique developmental tasks who lack protections from involuntary or coercive treatment, and whose parents and guardians need accurate information to make informed decisions about behavioral health treatment.
3. There is a lack of published research on efforts to change sexual orientation among children and adolescents; no existing research supports that mental health and behavioral interventions with children and adolescents alter sexual orientation. Given the research on the secondary outcomes of such efforts, the potential for risk of harm suggests the need for other models of behavioral health treatment.
4. Behavioral health professionals provide accurate information on sexual orientation, gender identity, and expression; increase family and school support; and, reduce rejection of sexual minority youth. Behavioral health practitioners identify sources of distress and work to reduce distress experienced by children and adolescents. Behavioral health professionals provide efforts to encourage identity exploration and integration, adaptive coping, and family acceptance to improve psychological well-being.

Professional Consensus on Gender Identity and Gender Expression in Youth

Consensus on the Overall Phenomena of Gender Identity and Gender Expression

1. Variations in gender identity and expression are normal aspects of human diversity and do not constitute a mental disorder. Binary definitions of gender may not reflect emerging gender identities.
2. Pre-pubertal children and peri-pubertal adolescents who present with diverse gender expressions or gender dysphoria may or may not develop a transgender identity in adolescence or adulthood. In pubertal and post-pubertal adolescents, diverse gender expressions and transgender identity usually continue into adulthood.

Consensus on Efforts to Change Gender Identity

3. There is a lack of published research on efforts to change gender identity among children and adolescents; no existing research supports that mental health and behavioral interventions with children and adolescents alter gender identity.
4. It is clinically inappropriate for behavioral health professionals to have a prescriptive goal related to gender identity, gender expression, or sexual orientation for the ultimate developmental outcome of a child's or adolescent's gender identity or gender expression.
5. Mental health and behavioral interventions aimed at achieving a fixed outcome, such as gender conformity, including those aimed at changing gender identity or gender expression, are coercive, can be harmful, and should not be part of treatment. Directing the child or adolescent to conform to any particular gender expression or identity, or directing parents and guardians to place pressure on the child or adolescent to conform to specific gender expressions and/or identities, is inappropriate and reinforces harmful gender stereotypes.

Consensus on Appropriate Therapeutic Intervention for Youth with Gender-Related Concerns

6. Children and adolescents experiencing gender-related concerns are an especially vulnerable population with unique developmental tasks. Parents and guardians need accurate scientific information to make informed decisions about appropriate mental health and behavioral interventions, including whether or not to initiate a social gender transition or, in the case of peri-pubertal, pubertal, and post-pubertal adolescents, medical intervention. Treatment discussions should respect the child's and adolescent's developing autonomy, recognizing that adolescents are still transitioning into adult decision-making capacities.
7. Approaches that focus on developmentally-appropriate identity exploration, integration, the reduction of distress, adaptive coping, and family acceptance to improve psychological well-being are recommended for children and adolescents of all ages experiencing gender-related concerns.

Pre-Pubertal Children

8. Gender expression and gender identity are interrelated and difficult to differentiate in pre-pubertal children, and are aspects of identity that develop throughout childhood. Therefore, a detailed psychological assessment should be offered to children and families to better understand the present status of a child's gender identity and gender expression, as well as any associated distress.

Peri-Pubertal Adolescents

9. For peri-pubertal adolescents, the purpose of pubertal suppression is to provide time to support identity exploration, to alleviate or avoid potential distress associated with physical maturation and secondary sex characteristics⁸, and to improve future healthy adjustment. If pubertal suppression is being considered, it is strongly recommended that parents or guardians and medical providers obtain an assessment by a licensed behavioral health provider to understand the present status of a peri-pubertal adolescent's gender identity or gender expression and associated distress, as well as to provide developmentally-appropriate information to the peri-pubertal adolescent, parents or guardians, and other health care professionals involved in the peri-pubertal adolescent's care. The purpose of the assessment is to advise and inform treatment decisions regarding pubertal suppression after sharing details of the potential risks, benefits, and implications of pubertal suppression, including the effects of pubertal suppression on behavioral health disorders, cognitive and emotional development, and future physical and sexual health.

Pubertal and Post-Pubertal Adolescents

10. Decision-making regarding one's developing gender identity is a highly individualized process and takes many forms. For pubertal and post-pubertal adolescents, if physical gender transition (such as hormone therapy or gender affirming surgeries) is being considered, it is strongly recommended that adolescents, parents, and providers obtain an assessment by a licensed behavioral health provider to understand the present status of an adolescent's gender identity and gender expression and associated distress, as well as to provide developmentally-appropriate information to adolescents, parents or guardians, and other health care professionals involved in the pubertal or post-pubertal adolescent's care. If physical transition is indicated, the potential risks, benefits, and implications of the transition-related procedures being considered – including the effects on behavioral health disorders, cognitive and emotional development, and potentially irreversible effects on physical health, fertility, and sexual health – are presented to the adolescent and parents or guardians.

Withholding timely physical gender transition interventions for pubertal and post-pubertal adolescents, when such interventions are clinically indicated, prolongs gender dysphoria and exacerbates emotional distress.